

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, 15/64, 15/67, 15/79	A1	(11) International Publication Number: WO 98/37189 (43) International Publication Date: 27 August 1998 (27.08.98)
(21) International Application Number: PCT/US98/03699 (22) International Filing Date: 25 February 1998 (25.02.98) (30) Priority Data: 60/038,500 25 February 1997 (25.02.97) US (71) Applicants (for all designated States except US): QBI ENTERPRISES LTD. [IL/IL]; Weizmann Scientific Park, P.O. Box 741, 74106 Nes-Ziona (IL). YISSUM RESEARCH DEVELOPMENT COMPANY [IL/IL]; Hebrew University of Jerusalem, Jabotinsky Street 46, 91042 Jerusalem (IL). (71) Applicant (for MW only): KOHN, Kenneth, I. [US/US]; 6761 Alderley Way, West Bloomfield, MI 48322 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KESHET, Eli [IL/IL]; Rechov Even Shmuel 24, 97234 Jerusalem (IL). STEIN, Ilan [IL/IL]; Ein Karem Bet 32, 95742 Jerusalem (IL). ITIN, Ahuva [IL/IL]; Rechov Haportzin 25, 92541 Jerusalem (IL). (74) Agents: KOHN, Kenneth, I. et al.; Kohn & Associates, Suite 410, 30500 Northwestern Highway, Farmington Hills, MI 48334 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: IRES SEQUENCES WITH HIGH TRANSLATIONAL EFFICIENCY AND EXPRESSION VECTORS CONTAINING THE SEQUENCE (57) Abstract An isolated and cloned translation control element, and analogues thereof, having the nucleotide sequence as set forth in SEQ ID NO:7 and designated SP163, are disclosed. The translation control element controls cap-independent mRNA translation via an internal ribosome entry site (IRES). The present invention provides expression vectors comprising the translation control element SP163 or its analogues operatively linked to a gene sequence to be expressed. In alternative embodiments, the expression vector comprises at least two nucleic acid sequences to be translated and SP163 is operatively linked to at least one of the sequences to be translated. The sequences to be translated may be linked to only one promoter in an embodiment. The present invention further provides a method for facilitating and enhancing cap-independent translation of mRNA by including in an expression cassette a translation control element having the nucleotide sequence as set forth in SEQ ID NO:7 and designated SP163.		

Document 11
Cited in IDS for CELL017
Serial No. 09/814,357,
filed March 21, 2001

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

IRES SEQUENCES WITH HIGH TRANSLATIONAL EFFICIENCY
AND EXPRESSION VECTORS CONTAINING THE SEQUENCE

BACKGROUND OF THE INVENTION

5

1. FIELD OF THE INVENTION

The present invention relates to an isolated and cloned DNA sequence element that can be incorporated into expression vectors for the purpose of improving translation of a given mRNA and to enable the translation-
of the mRNA in a cap-independent manner.

2. DESCRIPTION OF RELATED ART

There is an extensive and growing need to produce eukaryotic gene products in eukaryotic cells. Methods are needed to maximize expression of a desired gene and production of the gene product in eukaryotic cells. Initiation of eukaryotic protein synthesis involves about ten initiation factors, eIFs. The eIF4 group of initiation factors collectively catalyze the recognition of the mRNA cap, the unwinding of mRNA secondary structure, and the binding of mRNA to the 43S preinitiation complex [Pain, 1996]. The selection of a particular mRNA from the pool of translatable RNAs is determined by the relative efficiency of translation initiation by ribosome scanning and is largely governed by the composition and structure of the 5'-untranslated region (5'UTR) of the mRNA [Kozak, 1991; Sonenberg, 1996].

In certain instances translation of the transcribed, processed, and end-modified (i.e. capped and polyadenylated) mRNA is the limiting step in the production of the protein [Mathews et al., 1996; Meyuhas et al., 1996]. Translation is initiated by mRNA-protein interactions preceding the engagement of the small

ribosomal subunit (40S) with the mRNA. Thus, limited availability of cap-binding proteins and competition with other cellular mRNAs for that proteins can be a rate-limiting factor in translation of the desired protein.

5 This limitation can be even more pronounced under stress conditions (e.g. heat shock, hypoxia, nutrient deprivation) in which cap-dependent translation is markedly compromised. [Mathews et al., 1996; Meyuhas et al., 1996]

10 An alternative mode of translation is one in which ribosomes bind to the mRNA independent of the cap structure using an internal ribosome entry site (IRES), a specialized sequence within the 5' untranslated regions that directly promote ribosome binding, independent of a
15 cap structure. IRES elements were first discovered in picornaviral mRNAs which are naturally uncapped but nonetheless efficiently translated [Jang et al., 1988; Pelletier and Sonenberg, 1988; Oh and Sarnow, 1995]. Subsequently, it was found that some cellular RNAs which
20 are normally capped can be translated either by the 5' end-dependent scanning mechanism or by an internal ribosome binding mechanism. Generally, IRES cannot be identified by sequence homology; known IRES have been identified and defined functionally [Mountford and Smith,
25 1995]. It appears that it is the conformation of the IRES sequence that enables the binding on the ribosome.

The list of cellular genes shown to contain sequences mediating internal initiation within their 5'UTR includes the immunoglobulin heavy chain binding
30 protein (BiP) [Macjak and Sarnow, 1991], antennapedia [OH et al., 1992], fibroblast growth factor (FGF) [Vagner et al., 1995], platelet-derived growth factor-B (PDGF-B) [Bernstein et al., 1997], insulin-like growth factor II (IGF-II) [Teerink et al., 1995], and the translation
35 initiation factor eIF4G [Gan and Rhoads, 1996]. The

potential utility of a cap-independent translation mode is best demonstrated in the case of viral RNAs in circumstances where cap-dependent translation is completely abrogated (through cleavage of an essential cap-binding protein by a virus-encoded protease) and the translation machinery is taken-over by IRES-containing viral RNA [Pelletier and Sonenberg, 1988]. The option of internal initiation is an advantage for competition with other mRNAs when certain components of the eIF4 complex become rate-limiting and this option provides a given mRNA the ability to be translated at times when cap-dependent translation is compromised. Such circumstances may develop under hypoxia where overall protein synthesis is significantly inhibited [Heacock and Sutherland, 1988; Kraggerud et al., 1995] or other stress conditions.

It would be useful to have a small sequence element, derived from a naturally-occurring cellular 5'UTR, that endows any desired gene with the ability to be more efficiently translated, in general, and to be translated in a cap-independent manner, in particular. That is, it would be useful to have additional IRES sequences with high translational efficiency, to use in expression vectors, to control mRNA translation and therefore protein synthesis as well as in gene therapy vectors.

SUMMARY OF THE INVENTION

According to the present invention, an isolated and cloned translation control element, and analogues thereof, having the nucleotide sequence as set forth in SEQ ID No:7 and designated SP163, is disclosed. The translation control element controls cap-independent mRNA translation via an internal ribosome entry site (IRES). The present invention provides expression vectors comprising the translation control element SP163 or its analogues operatively linked to a gene sequence to be

expressed. In alternative embodiments, the expression vector comprises at least two nucleic acid sequences to be translated and SP163 is operatively linked to at least one of the sequences to be translated. The sequences to be translated may be linked to only one promoter in an embodiment.

The present invention provides a method for facilitating and enhancing cap-independent translation of mRNA by including in an expression cassette a translation control element having the nucleotide sequence as set forth in SEQ ID No:7 and designated SP163.

The novel sequence element (designated SP163) is composed of sequences derived from the 5'-UTR of VEGF (Vascular Endothelial Growth Factor gene), however, in a novel arrangement that was presumably generated through a previously unknown mode of alternative splicing. Functional analysis has shown that SP163 functions as a significant stimulator of translation and as a mediator of cap-independent translation. Interestingly, the full-length 5'-UTR of VEGF has fair IRES activity indicating that these stronger activities were generated by a specific molecular event causing the juxtaposition of two specific 5'-UTR segments.

The advantages of SP163 is that it is a natural cellular IRES element with a superior performance as a translation stimulator and as a mediator of cap-independent translation relative to known cellular IRES elements. Another advantage of SP163 is that these functions are maintained under stress conditions.

DESCRIPTION OF THE DRAWINGS

Other advantages of the present invention will be readily appreciated as the same becomes better understood by reference to the following detailed description when

considered in connection with the accompanying drawings wherein:

FIGURES 1A-B are schematic diagrams of vectors and constructs used in this study. Figure 1A is bicistronic expression vector in which expression of a bicistronic mRNA is driven by a CMV virus promoter. A firefly luciferase (LUC) is translated from the first cistron and a secreted alkaline phosphatase (SeAP) from the second cistron. Putative IRES elements are inserted into the intercistronic space (ICS). Figure 1B is Monocistronic expression vector in which SeAP expression is driven by a CMV virus promoter. Sequence elements tested for a translation- modulating activity are inserted upstream of the SeAP coding region.

FIGURE 2A-C are graphs showing the production of Luciferase and SeAP from a bicistronic mRNA in stably transfected C6 cells wherein Figure 2A is a graph of production of SeAP from the downstream cistron. Pools of stably transfected C6 clones were grown to 70% confluence, medium was replaced with a fresh medium (t=0) and aliquots were withdrawn at the indicated time points and analyzed for SeAP activity. Activity is expressed as cumulative units of SeAP in 1 ml of medium and is standardized to total protein. For testing SeAP production under hypoxia, cells were shifted to 1% oxygen at t=0 and further analyzed as above. Figure 2B is a bar graph showing ratio of SeAP/LUC production in the different transfectants. SeAP/LUC ratio was calculated from the respective activities determined at the endpoint of the experiment (t=24hrs.). Figure 2C is a bar graph showing that under hypoxia stress where translation of LUC gene from the upstream cistron was reduced the rate of SeAP production was unaffected indicating internal initiation. Plasmid designation is as shown in Figure 1. N=Normoxia, H=Hypoxia.

FIGURE 3A-B are photographs of Northern blot analysis of mRNAs transcribed from bicistronic plasmids. RNA was extracted from untransfected C6 cells and from the same stably transfected C6 pools used in the experiment shown in Figure 2. 20 μ g of each RNA was electrophoresed in two parallel lanes, blotted and hybridized with either a LUC-specific probe (A) or a SeAP-specific probe (B). To assure equal loading, ribosomal RNAs were stained with methylene blue prior to hybridization. Lane 1 - untransfected C6 cells; Lane 2 - B/0; Lane 3 - B/UTR; Lane 4 - B/Bip.

FIGURE 4 is a photograph of a Northern blot analysis of VEGF mRNA species expressed under conditions of severe hypoxia. C6 cells were grown under normoxia (N) or were exposed to 1% oxygen for 4 hours (4H) or 16 hours (16H). 5 μ g of poly (A)⁺ RNA from each culture was subjected to a Northern blot analysis using a VEGF-specific probe. The bold arrow points at the most frequently encountered 3.7 kB VEGF mRNA.

FIGURE 5 is a graph of IRES activity of SP163. 293 cells were transiently transfected with each of the indicated bicistronic plasmids and SeAP and LUC activities were determined 36 h post-transfection as described under 'Methods'. Results are expressed as a SeAP/LUC ratio and represent the average of 4 independent transfections for each plasmid. Plasmid designations are as shown in Fig. 1. B/ Δ SP163 is a deletion mutant of SP163 missing the first 31 nucleotides. B/SP163/M5' is a substitution mutation wherein the 9 nucleotides from the 5' terminus of SP163 are substituted. B/SP163/M3' is a substitution mutation of the 5 nucleotides from the 3' terminus. Inset is a photograph of a Northern blot analysis with a SeAP-specific probe and a LUC-specific probe performed on 20 μ g of the total RNA extracted from transfected 293 cells 36 h post-transfection. Lane 1 -

mock-transfected cells; lane 2 - B/0-transfected cells;
lane 3 - B/UTR-transfected cells; lane 4 - B/SP163-
transfected cells.

FIGURE 6 is a graph showing the effectiveness of
5 SP163 when compared to another cellular IRES, that of the
grp78 gene.

FIGURE 7 is a graph showing results obtained for two
representative species, rat C6 cells and human Hela
cells.

10 FIGURE 8A-B are graphs showing that the enhancing
performance of SP163 is retained when the expression
vector is stably integrated into the host cell genome.

FIGURE 9 is a graph showing the relative stimulation
of SeAP activity and SeAP activity in pBKC/163S was 4
15 fold higher than pCIBB/S and over 6-fold higher than
pCImc/S.

FIGURE 10A-C are photographs of gel electrophoresis
showing Northern blot analysis of the amounts of mRNA of
the reporter genes (SeAP and Luc) that are transcribed
20 from the different vectors.

FIGURE 11 is a graph showing translation enhancing
activity of VEGF 5'UTR sequences under normal (N) and
hypoxia (H) conditions. C6 cells were stably transfected
with the monocistronic constructs indicated. SeAP
25 activity was analyzed in pools of transfected clones as
described under 'Methods' and values were standardized to
total protein. For analysis of SeAP activity under
hypoxia, cells were shifted to 1% oxygen 24 hours before
sampling. Inset is a photograph of Northern blot
30 analysis of SeAP-containing mRNAs in stably-transfected
pools. Lane 1-M/0-transfected cells; lane 2 - M/UTR-
transfected cells; lane 3 - M/SP163-transfected cells;
lane 4 - non-transfected cells.

DETAILED DESCRIPTION OF THE INVENTION

According to the present invention, an isolated and cloned translation control element, and analogues thereof, having the nucleotide sequence as set forth in SEQ ID No:7 and designated SP163, is disclosed. The translation control element controls cap-independent mRNA translation via an internal ribosome entry site (IRES).

The term Analogue as used herein is defined as a nucleotide sequence variant (alternatively the terms alteration, sequence alteration, sequence variant can be used) with some differences in their nucleotide sequences as compared to the native sequence of SEQ ID No:7 but, such that the conformational folding of the sequence, to allow its use as an IRES, is not compromised. Ordinarily, the analogue will be generally at least 70% homologous over any portion that is functionally relevant. In more preferred embodiments the homology will be at least 80% and can approach 95% homology to the nucleotide sequence. The nucleotide sequence of an analogue may differ from that of the translation control element of the present invention when at least one nucleotide is deleted, inserted or substituted, but the translation sequence element remains functional as an IRES. Functionally relevant refers to the biological property of the sequence and in this context means a conformational or other aspects of the molecule that allow functioning as an enhanced IRES as described herein.

The present invention provides expression vectors comprising the translation control element, designated SP163, operatively linked to a gene sequence to be expressed. In alternative embodiments the expression vector comprises at least two nucleic acid sequences to be translated and the translation control element is

operatively linked to at least one of the sequences to be translated. Vectors are known or can be constructed by those skilled in the art and contain all expression elements necessary to achieve the desired transcription of the sequences in addition to the sequence of the present invention as shown in the Examples herein below. The vectors contain elements for use in either procaryotic or eucaryotic host systems depending on their use. One of ordinary skill in the art will know which host systems are compatible with a particular vector.

Additional features can be added to the vector to ensure its safety and/or enhance its therapeutic efficacy as known in the art. Such features include, for example, markers that can be used to negatively select against cells infected with the recombinant virus. An example of such a negative selection marker is the TK gene that confers sensitivity to the anti-viral drug gancyclovir. Negative selection is therefore a means by which infection can be controlled because it provides inducible suicide through the addition of antibiotic. Such protection ensures that if, for example, mutations arise that produce altered forms of the viral vector or sequence, cellular transformation will not occur. Features that limit expression to particular cell types can also be included. Such features include, for example, promoter and regulatory elements that are specific for the desired cell type in addition to the translation control sequence of the present invention. As shown in the Examples herein the sequences can be used in all cell types in both vectors for use in cell culture and in vivo gene therapy.

The present invention also provides a method for facilitating preferential translation of a gene of interest over the bulk of cellular mRNAs which are not of interest. In other words allowing restriction of protein

production in the host cell to essentially the protein/peptide coded for by the gene/nucleotide sequence of interest. The method provides the steps of including in an expression vector a translation control sequence of the present invention as set forth in SEQ ID No:7, 5 operatively linked to the gene of interest. The vector is then expressed in host cells as is known in the art. The cells are then treated with compounds that inhibit cap-dependent translation. The reagents that are 10 administered to the cell can include, for example, a virus-derived protease or anysomycin or other agents.

The present invention provides a second method for facilitating preferential translation of a gene of interest over the bulk of cellular mRNAs. In this 15 further embodiment, the steps provide constructing an expression vector including a translation control sequence of the present invention as set forth in SEQ ID No:7, operatively linked to the gene of interest as well as any other required regulatory elements required for 20 the sequence of interest. The vector is then expressed in host cells, as is known in the art. The host cells are then cultured under conditions which will induce cellular stress such as hypoxia, heat shock, hypoxia, hypoglycemia, iron deprivation and other 25 conditions/treatments/compounds which will induce cellular stress such that cap-independent mRNA translation is active and cap-dependent translation is impaired.

As shown in the Examples, with the aid of reverse 30 transcription-PCR, the use of mouse cytoplasmic mRNA as a template, and the primers indicated in Table 1 (corresponding to the 5' and 3' boundaries of the 5'-untranslated region of VEGF), the sequence of the present invention shown in Table 1, SEQ ID No:7, was amplified 35 and subsequently cloned.

Comparison of this sequence (designated SP163; SEQ ID No:7) with known 5'-UTR VEGF sequences from rat [Levy et al, 1995], mouse [Shima et al, 1995] and human [Tischer et al, 1991] revealed that it is composed of the first 31 nucleotides of the 5'-UTR (including the cap) adjoined to the last 132 nucleotides of the 5'UTR (i.e. up to the initiator ATG condon). It appears that SP163 is generated by a splicing event that removes the bulk (>800 nucleotides) of internal 5'-UTR sequences.

Sequencewise, there is no obvious resemblance among the known cellular IRESSs, nor do any of the cellular IRESSs show a significant homology with picornavirus IRESSs. For example, none of the known cellular IRESSs has a pyrimidine-rich tract located 25 nucleotides upstream of the initiation site, as in the picornavirus IRESSs [Ehrenfeld, 1996]. VEGF share a considerable sequence homology with another cellular gene that has an IRES element namely, PDGF. However, a search for sequence homology between the respective 5'UTRs have shown that, despite a similarity in length and overall high GC content, there is no significant homology in primary sequences.

Further Examples demonstrate that SP163 significantly improves production of a given protein, when included in various contexts of both monocistronic and bicistronic expression vectors. SP163, was introduced into the basic mono-cistronic expression vector pCIbb, and into the new bi-cistronic expression vectors pBIC-LS to create improved vectors (Figures 1A,B; Table 2). Experiments describing the performance of SP163 in the two vector systems are summarized as follows and detailed in the Examples.

In the mono-cistronic vector type (pBKC; M/O), Figure 1B, SP 163 is introduced between the strong CMV promoter and the target gene. In this configuration, it

enhances the expression of the gene located downstream. The secreted alkaline phosphatase reporter (SeAP) system is used to measure SP163 activity for testing its performance in different cellular conditions. This
5 reporter system is especially convenient since it monitors the reporter activity in the growth media of the cells and allows the performance of elaborate kinetics studies.

In the bi-cistronic vector system, Figure 1A, SP163
10 is introduced between the two target recombinant genes.— The expression of the first gene is driven by the strong CMV promoter, while that of the second is governed by cap-independent translation via the IRES element SP163. In the reporter version, bi-cistronic vector pBIC/LS
15 (B/O; Figure 1A), SP163 or any other IRES fragments studied are cloned into the unique multiple-cloning site element downstream to the luciferase (Luc) gene. The IRES element directs the expression of Secreted Alkaline Phosphatase (SeAP) via a cap-independent mechanism. The
20 activity of the luciferase gene is used to normalize changes in transcription between different vectors containing different IRESs.

This embodiment of pBIC with the IRES of the present invention will allow the cloning of any two target genes,
25 such as recombinant genes I and II, so that gene I will be expressed via the CMV promoter and gene II will be expressed via SP163.

Transient transfection is commonly used for characterization of the function of a cloned gene. In
30 order to analyze the ability of SP163 to stimulate expression in transient transfection, the vectors pBKCS and pBKCS163S were introduced into different cell lines by liposomes mediated transfection. As shown in the Examples, the results obtained indicate that in all the
35 species tested the addition of SP163 to the pBKCS vector

leads to a significant stimulation of expression of 5-10 fold as measured by the reporter SeAP.

5 In the case of VEGF, internal ribosome entry circumvents the need for a troublesome ribosome scanning through the exceptionally long, highly structured 5'UTR which could have rendered translation an inefficient process [Kozak, 1991]. Among its many functions, VEGF plays an important role in maintaining vascular homeostasis and controlling vascular permeability [Senger
10 et al., 1993]. Notably, its unscheduled downregulation — at critical developmental timings may cause pathologic regression of existing vessels [Alon et al., 1995]. Thus, the capacity for cap-independent translation may serve as a safeguard, preventing the deleterious
15 consequence of VEGF under-translation in circumstances where cap-dependent translation is transiently compromised.

The above discussion provides a factual basis for the use of IRES sequences with high translational
20 efficiency and expression vectors containing the sequence. The methods used with and the utility of the present invention can be shown by the following non-limiting examples and accompanying figures.

25

EXAMPLES

GENERAL METHODS:

General methods in molecular biology: Standard molecular biology techniques known in the art and not specifically described are generally followed as in
30 Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory, New York (1989, 1992), and in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Maryland (1989). Polymerase chain reaction (PCR) is carried out generally
35 as in *PCR Protocols: A Guide To Methods And Applications*,

Academic Press, San Diego, CA (1990). Reactions and manipulations involving other nucleic acid techniques, unless stated otherwise, are performed as generally described in Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, and methodology as set forth in United States patents 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057 and incorporated herein by reference. *In-situ* (In-cell) PCR in combination with Flow Cytometry can be used for detection of cells containing specific DNA and mRNA sequences [Testoni et al, 1996, Blood 87:3822]. The vectors of the present invention are synthesized as described herein and by any method known in the art.

Construction of bicistronic and monocistronic expression plasmids. The basic bicistronic vector (designated B/0) and basic monocistronic vector (designated M/0) used in this study for insertion of 5'UTR elements were provided by QBI Enterprises Inc. (Rehovot, Israel). Their structures are described in Figs. 1A&B.

5'UTR elements were obtained by RT-PCR amplification using the specific primers indicated below. Primers were designed in a way that each amplified fragment was bounded by *Xho*I (5') and *Nco*I (3') sites and was inserted between the *Xho*I (5') and *Bsm*BI (3') sites of the intercistronic spacer of the bicistronic vector B/0 or into the same sites of the monocistronic vector M/0. In either case, the hybrid *Nco*I/*Bsm*BI site recreated the initiator ATG codon of the SeAP cistron.

Cloning VEGF and BiP 5'UTRs: RNAs from hypoxic NIH3T3 cells or 293 cells, respectively, were reverse transcribed using 10 µg of total RNA. PCR amplification was carried out using a Taq DNA Polymerase possessing a proofreading activity (Pwo DNA Polymerase, BOEHRINGER) and the following oligonucleotide primers: VEGF 5'

primer: 5' CTCGAGCGCAGAGGCTTGGGGC (SEQ ID No:1). VEGF
3' primer: 5'CCATGGTTTCGGAGGCCGTCCG 3' (SEQ ID No:2)
corresponding to nucleotides 1218-1235 and 2234-2214,
respectively, of the mouse VEGF gene (GeneBank accession
5 U41383). The oligonucleotide primers used to amplify the
full length VEGF 5'UTR were also used to amplify the new
splice variant of VEGF 5'UTR designated SP163 which was
also inserted into the B/0 and M/0 vectors in the same
way as described above. BiP 5' primer:
10 5'CTCGAGAGGTCGACGCCGCGCCAAGACA (SEQ ID No:3). BiP 3'
primer: 5'CCATGGCTTGCCAGCCAGTTGGGCAGC (SEQ ID No:4)
(corresponding to nucleotides 372-392 and 592-572,
respectively, of the human BiP gene, Genebank accession
M19645).

15 All plasmid constructs were sequenced to verify
their structure and the preservation of open reading
frames within PCR amplified fragments. A summary of
bicistronic and monocistronic constructs used in this
study is shown in Table 2 and Fig. 1A-B..

20 Cell culture and DNA transfections: C6 cells, a
clonal glial cell line derived from a rat glial tumor
[Benda et al., 1968], were grown in Dulbeco-Modified
Eagle's medium (DMEM) containing 5% fetal calf serum and
antibiotics. The human cell line 293 (adenovirus-
25 transformed fetal kidney cells), mouse NIH/3T3 cells and
rat primary astrocytes were maintained in Dulbeco-
Modified Eagle's medium (DMEM) containing 10% fetal calf
serum and antibiotics. Primary rat astrocytes were
prepared as described previously [Frangakis and
30 Kimelberg, 1984].

Where indicated, cells were exposed to hypoxia (1%
oxygen) using a CO₂/O₂ incubator (Forma Scientific) in
which oxygen levels are monitored and adjusted
automatically.

Transfections were performed using Lipofectin (GIBCO BRL) as follows: Cells were plated at a density of $1-2 \times 10^5$ Cells per 60-mm plate and grown for 24 hours, medium was replaced with a serum-free DMEM and 200 μ l of DNA-Lipofectin suspension (containing 2 μ g supercoiled plasmid and 15 μ g of Lipofectin, preincubated at room temperature for 40 minutes) were added in a dropwise manner. Fourteen hours later, medium was replaced with a medium supplemented with serum and antibiotics and cells were incubated for additional 48 hours before splitting into a selection medium containing G418 (2mg/ml). Selection continued for 10-14 days and >50 individual G418-resistant clones were pooled for each plasmid.

For transient transfections, cells were seeded at a density of 2×10^5 cells per 1mm well and transfected 24 hours later with 1 μ g of DNA per well using the calcium phosphate procedure [Jordan et al., 1996]. Cells and media were harvested 36 hours post-transfection. Cell extracts were analyzed for luciferase activity or used for mRNA preparation (in the latter case, 10 μ g of DNA was used to transfect $2-2.5 \times 10^6$ cells per 100 mm plate), while the media was used for analysis of SeAP activity.

RNA analysis: Total RNA was prepared by the guanidinium thiocyanate/phenol-chloroform extraction procedure [Chirgwin et al., 1979]. mRNA was purified from total cellular RNA using the mRNA Separator Kit (CLONTECH).

Analysis of mRNA in polysomal fractions: Protein synthesis was instantaneously arrested by treating the culture with cycloheximide (90 μ g/ml) for 10 minutes. Cells were collected, washed with phosphate-buffered saline and kept at -70°C until analyzed. Cell lysis, size fractionation of polysomes by sedimentation through sucrose gradients, and preparation of ribosome-associated

RNAs were performed as described previously [Meyuhas et al., 1987].

Northern blotting: RNA was denatured in glyoxal and electrophoresed through a 1.0% agarose gel. RNAs were transferred onto a nylon-based membrane (GeneScreen plus, NEN) by the capillary blot procedure, and were hybridized with the indicated cDNAs, labeled with ³²P by randomly primed DNA synthesis. For standardization, blots were rehybridized with a β -actin probe or, alternatively, ribosomal RNAs were visualized by staining with methylene blue prior to hybridization. For relative quantification, autoradiograms were scanned using a PhosphorImager (FUJIX BAS 1000 Bio-Imaging Analyzer). Hybridization probes used were: VEGF (a 0.6 kB long mouse cDNA fragment encoding VEGF165) [Shweiki et al., 1992], β -actin [Minty et al., 1981], luciferase (a 1.7 kB long cDNA fragment including all of the coding region of firefly luciferase), and SeAP (a 1.5 kB long cDNA fragment including the coding region of the human SeAP).

RT-PCR: semi-quantitative RT-PCR was performed using a single tube, one step RT-PCR system (Access RT-PCR system, PROMEGA) and 20 cycles of amplification. Oligonucleotide primers used for reverse transcription and amplification of the various splicing variants of VEGF mRNA were:

5'GAGAGAATGAGCTTCCTACAG 3' (SEQ ID No:5) and 5' TCACCGCCTTGGCTTGTCACA 3' (SEQ ID No:6) (derived from the common fifth and eighth exons, respectively). PCR products were resolved by agarose gel electrophoresis, and detected by blot-hybridization using a VEGF-specific, ³²P-labeled cDNA probe.

Luciferase and SeAP analysis: Luciferase enzymatic activity in the cell extracts was determined using a commercial Luciferase Assay System (PROMEGA) and the

procedure recommended by the supplier. Light generated was measured using Lumac/3M BIOCOUNTER M2010-luminometer.

SeAP activity released into the growth medium was determined as follows: medium was heated for 5 minutes at 65°C and clarified by centrifugation. Aliquots were diluted in a SeAP buffer (1M Diethanolamine pH 9.8, 0.5 mM MgCl₂, 10m M L-homoarginine) in a 96 well plate and the enzymatic reaction (at 37°C) initiated with the addition of 20 µl of 120mM p-nitrophenylphosphate. The reaction product was determined using an ELISA reader at A450-630. The amounts of conditioned medium added and incubation times used were adjusted so that readings were within the linear range of the calibration curve obtained with a purified SeAP standard. Total protein was determined by the Bradford method [Bradford, 1976].

EXAMPLE 1

ANALYSIS OF THE FULL 5'UTR OF VEGF

The 5'UTR of vascular endothelial growth factor (VEGF) has several features which are incompatible with efficient ribosomal scanning. First, it is considerably longer (1014 nucleotides in the mouse) than most eukaryotic 5'UTRs. Second, it has a high G/C content (comprising 64% of 5'-UTR sequences and 80% of the 100 nucleotides preceding the translation initiation condon) and can potentially form complex stable secondary structures. Third, the 5'UTR contains a short open reading frame bounded by in-frame initiation and termination condons. The inherent difficulty for efficient ribosome scanning, on the one hand, and the possibility of IRES, on the other hand, prompted Applicants to examine if the VEGF mRNA is translated in a cap-independent mode.

In review, VEGF plays a key role in the initiation of blood vessel formation. VEGF is produced by and

secreted from the tissue towards which new blood vessels eventually extend. A large body of evidence supports the premise that the amount of VEGF produced determines the magnitude of the angiogenic response through interaction with cognate receptors expressed on nearby endothelial cells. VEGF expression is tightly regulated *in vivo* and a deviation from the normal levels of VEGF might be detrimental to the vasculature. For example, a reduction of VEGF gene dosage by one half (in mice heterozygous for an inactivating mutation in VEGF) leads to severe vascular defects and early embryonic lethality [Carmeliet et al., 1996; Ferrara et al., 1996]. Conversely, over-expression of VEGF may lead to pathologies like retinopathy resulting from excessive proliferation of blood vessels [Aiello et al., 1994; Pe'er et al., 1995].

VEGF expression can be modulated *in vitro* by a variety of agents, including a number of cytokines, growth factors and steroid hormones. Yet, from a physiological point of view, regulation of VEGF by hypoxia is the most significant. Inefficient vascular supply and the resultant reduction in tissue oxygen tension, often lead to compensatory neovascularization acting to satisfy the metabolic needs of the tissue. Hypoxia-induced VEGF was found to be the key mediator of this feedback response [Plate et al., 1992; Shweiki et al., 1992]. VEGF is regulated by hypoxia at both the transcriptional and post-transcriptional levels.

Transcriptional regulation of VEGF is mediated by the transcription factor, hypoxia-inducible factor 1 (HIF-1) which accumulates under conditions of hypoxia and activates VEGF transcription through binding to specific promoter sequences [Forsythe et al., 1996]. Hypoxia also leads to stabilization of VEGF mRNA [Ikeda et al., 1995; Shima et al., 1995; Stein et al., 1995]. The intrinsically short half-life of VEGF mRNA (approximately

30 minutes) is significantly extended under stress, presumably through hypoxia-augmented binding of yet unidentified protein(s) to its 3'untranslated region [Levy et al., 1996]. Both mechanisms act to increase steady-state levels of VEGF mRNA. However, it is not known whether production of VEGF is also regulated at the level mRNA translation and, in particular, whether the mode or extend of VEGF translation is effected by hypoxia. VEGF is expected to be maximally produced under hypoxia in order to fulfill its function as a mediator of hypoxia-driven angiogenesis. These considerations led to the examination of the performance of the 5'UTR under conditions of hypoxic stress.

To determine the efficiency by which VEGF mRNA is translated, primary astrocyte cultures were grown, cytoplasmic extracts prepared and fractionated by centrifugation through a sucrose gradient into subpolysomal and polysomal fractions. The relative abundance of VEGF mRNA in these fractions was determined by quantitative RT-PCR. The bulk of VEGF mRNA was associated with polyribosomes, indicating that the majority of VEGF mRNA is engaged in active protein synthesis. These results suggested that VEGF mRNA is efficiently translated under normal growth conditions despite its cumbersome 5'UTR.

To determine whether translation of VEGF mRNA is inhibited under conditions of severe hypoxia, a similar analysis was carried out using cultures grown for 16 hours under 1% oxygen. Astrocytes were chosen for this study since they are the first cells to respond to hypoxia of neuronal tissues by upregulating VEGF mRNA expression and are, therefore, the cells responsible for the feedback angiogenic response [Stone et al., 1995]. The fraction of RNA associated with polyribosomes was reduced by 50% in comparison with cells grown under

normoxia. This result is consistent with previous observations that hypoxia causes about 30-50% inhibition in overall protein synthesis [Heacock and Sutherland, 1988; Kraggerud et al., 1995]. Despite the general
5 impairment of protein synthesis, the majority of VEGF RNA remained associated with polysomes. Notably, all isoforms of VEGF mRNA remained associated with the polyribosomal fraction, including the shortest VEGF121 which can accommodate no more than 3-4 ribosomes when
10 translated at a theoretical maximal rate. In contrast to VEGF, the fraction of mRNA encoding ribosomal protein L19 that was associated with polyribosomes was reduced 2-fold under hypoxia in comparison to normoxia. These results suggest that translation of VEGF mRNA is not a rate-
15 limiting step in production of this vital protein. Importantly, translation does not appear to be rate-limiting even under hypoxia where the steady-state level of VEGF mRNA is elevated by one order of magnitude [Shweiki et al., 1992].

20 The fact that translation of VEGF mRNA is very efficient despite its 'inappropriate' 5'UTR, prompted examination of whether VEGF mRNA can be translated by internal ribosome entry. Bicistronic mRNAs have been effectively used *in vivo* to demonstrate the existence of
25 IRES in both picornaviral [Pelletier and Sonenberg, 1988] and cellular [Gan and Rhoads, 1996; Macejak and Sarnow, 1991] mRNAs as discussed herein above. A eukaryotic expression vector was constructed in which a CMV virus promoter drives expression of a bicistronic RNA. The
30 first cistron in the bicistronic mRNA used in this study, encoded firefly luciferase (LUC) and should be translated by a conventional cap-dependent scanning mechanism. As ribosomes fail to continue scanning through the intercistronic region, the second cistron, encoding a
35 secreted alkaline phosphatase (SeAP), should be

translated only if preceded by an IRES. SeAP was chosen as the downstream cistron since secretion of the enzyme into the culture medium allows accurate quantification and continuous monitoring of protein synthesis.

5 To prevent leaky translation into the second
cistron, the intercistronic region included termination
condons in all possible reading frames. The 5'UTR of
VEGF was sub-cloned by precise PCR amplification,
10 inserted into the intercistronic spacer (ICS) region and
transfected into a C6 rat glioma cell line. Pools of
stably-transfected clones were prepared, each composed of
>50 individual clones. Analysis of pools assured that
possible differences in gene expression due to different
integration sites are averaged. Reporter gene activity
15 was determined in transfectants obtained with the
bicistronic vector alone (designated B/0) and with the
bicistronic vector containing the VEGF 5'UTR (designated
B/UTR). The two constructs expressed a bicistronic mRNA
from which LUC was translated with a comparable
20 efficiency. Analysis of the downstream cistron, however,
revealed that B/0 produced a negligible amount SeAP
activity, confirming a minimal, if any, readthrough of
ribosomes from the LUC to the SeAP cistron. In contrast,
B/UTR directed the production of high levels of SeAP in a
25 continuous manner (Fig. 3A), indicating that the 5'-
untranslated region of VEGF contains a functional IRES
element.

 To appreciate the relative strength of the VEGF
IRES, a comparison was made with the well-characterized
30 cellular IRES contained in the 5'UTR of BiP mRNA
[Pelletier and Sonenberg, 1988]. To this end, the 5'UTR
of BiP was inserted into the same site of the bicistronic
vector, pools of stable transfectants were prepared and
similarly analyzed for LUC and SeAP production. As shown

in Fig. 3B, the VEGF IRES was 5-fold more efficient than the BiP IRES in directing SeAP production.

While there is a clear rationale for the use of internal initiation in viruses, the advantage of internal initiation in cellular mRNAs is not always clear. In the case of VEGF, however, the capacity for cap-independent translation might be particularly advantageous in hypoxia where overall protein synthesis is compromised. Therefore, we have measured SeAP production, presumably by internal initiation, was unaffected by hypoxia. As shown in Figure 2A and 2C, the rate of SeAP production, presumably by internal initiation, was unaffected by hypoxia.

To rule out that in B/UTR-transfected cells the protein encoded by the second cistron is translated from a monocistronic SeAP mRNA, a RNA blot-hybridization analysis was carried out using both a LUC-specific probe and a SeAP-specific probe. A single band of RNA was detected in each case, corresponding to the expected size of the respective bicistronic transcript (the larger size of hybridized mRNA in B/UTR-transfectants is due to the addition of 1014 base pairs of 5'UTR sequence) (Figure 3A,B). Hybridization with both a LUC-specific probe and a SeAP-specific probes detected the same mRNA species, indicating that both cistrons are contained in a single transcripts. Importantly, the failure to detect a band corresponding in size to that of a presumptive monocistronic seAP mRNA ruled out the possibility that a significant amount of SeAP is translated from a monocistronic SeAP mRNA. Also note that the inclusion of the 5'UTR had no effect on the steady-state level of the bicistronic mRNA expressed, indicating that differences in measured LUC and SeAP activities are solely due to differences in translation efficiencies.

35

EXAMPLE 2

IDENTIFICATION OF SP163

Alternative splicing of VEGF RNA results in formation of four mRNA species encoding different isoforms of the protein [Ferrara et al., 1991]. These mRNAs differ with respect to the presence or absence of coding exons 6 and 7 and are coordinately regulated by hypoxia (i.e. production of all isoforms is equally increased by hypoxia) [Banai et al., 1994; Minchenko et al., 1994]. There is no evidence, however, for the existence of variant forms of VEGF mRNA which differ with respect to the 5'-3'-untranslated regions. Yet, when cells are grown under conditions of severe ischemia, additional new forms of VEGF mRNA accumulate, the size of which can not be accounted for by changes in the coding region alone (Fig. 4). Interestingly, smaller VEGF mRNAs have been encountered before but have generally been ignored [e.g. Mazure, 1996; Finkenzeller, 1995]. The appearance of these smaller mRNAs suggested that they might represent transcripts with a truncated 5'UTR, or a truncated 3'UTR, or both.

To examine the possibility that splice variants of the 5'UTR exist, mRNA from hypoxic NIH 3T3 cells was amplified by RT-PCR using oligonucleotide primers derived from the most distal and most proximal segments of the full length 5'UTR. The use of this primer combination assured that only 5'UTR sequences are amplified and only of transcripts sharing the same 5' terminus. Several fragments were amplified and verified by blot-hybridization as containing VEGF 5'UTR sequences. However, only one amplified fragment was cloned and further analyzed. The 163 nucleotides long sequence of this modified 5'UTR, designated SP163 (SEQ ID No:7), is shown in Table 1. Alignment of this sequence with that of the full length 5'UTR suggested that it was generated

by a splicing event that juxtaposed the 5' cap-containing segment of 31/32 nucleotide next to the 132/131 nucleotides preceding the initiator AUG codon. The experiments described below demonstrate the performance of the SP163 element as a mediator of cap-independent translation and its activity as a translation enhancer.

EXAMPLE 3

SP163 FUNCTION AS IRES

As described in Examples 1 and 2, SP163 was identified during the search for possible IRES sequences in the 5'UTR of VEGF. A series of further experiments were designed to characterize its activity as an IRES. The ability of SP163 to promote expression that is derived from the ribosomal entry site was tested in the bi-cistronic vector pBIC-LS. pBIC-LS contains luciferase gene as the first cistron and SeAP as the second cistron, in between is a multiple cloning site that allows the introduction of potential IRES sequences. The IRES activity of SP163 was compared to the parental vector pBIC/LS and to the entire 5'UTR of VEGF from which it was derived. It has stronger activity than the entire 5'UTR. The three plasmid pBIC/LS, pBIC/L5'UTRS and pBIC/L163S were introduced transiently into C6 glioma cells. SeAP activity reflects the IRES effectiveness, and luciferase activity is used to normalize in the overall changes in transcription. SP163 was generally over 5-8 fold more active as an IRES than the entire 5'UTR.

The SP163 element was inserted into the ICS region of the same bicistronic vector used to analyze the 'conventional' 5'UTR (Example 1). This plasmid, designated B/SP163 was transfected into the human 293 cell line alongside with plasmid B/0 (serving as a negative control) and plasmid B/UTR (for comparison of relative IRES strength). Proteins encoded by the first

and second cistrons were quantified following a transient transfection (Figure 5). Again, only negligible SeAP activity could be detected in cells transfected with B/0, confirming a non-leaky ribosome scanning through the first cistron. Consistent with the results of the stable transfections shown in Figure 2A-B, the full-length 5'UTR directed a considerable level of SeAP production also in cells of human origin. Remarkably and unexpectedly, SP163 was 5-fold more efficient than the full-length 5'UTR as IRES element.

SP163 also functions as an IRES in the CHO cell. pBIC-LS contains neomycin genes that allow selection following introduction and establishment of the vector into cells. Plasmid pBIC/LS (parental) and pBIC/L163S were transfected into the CHO cell. In this experiment, the effectiveness of SP163 was compared to another cellular IRES, that of the grp78 gene. The result of this experiment, presented in Figure 6, shows that SP163 function as an IRES after its establishment in the host genome. Its activity is over 25-fold over the background (pBIC/LS). Moreover, it is three-fold more effective as an IRES than the IRES of grp78, the best characterized cellular IRES known to date.

RNA blot-hybridization analysis was performed to assure that SP163-directed translation of SeAP was by internal ribosome entry. As shown in Figure 5 (inset), a single mRNA species, corresponding in size to the expected size of a bicistronic B/SP163 mRNA co-hybridized with the LUC-specific probe and the SeAP-specific probe and no monocistronic SeAP mRNA could be detected. This result indicated that SeAP was translated from a bicistronic RNA using a SP163-directed internal ribosome entry.

EXAMPLE 4

ACTIVITY OF SP163

Activity in transient expression in cell lines of different species. Transient transfection is commonly used for characterization of the function of a cloned gene. In order to analyze the ability of SP163 to stimulate expression in transient transfection, 5 µg of plasmid DNA of the vectors pBKCS and pBKC163S was introduced into different cell lines by liposomes mediated transfection. Twenty four hours after the transfection, the growth media was changed and SeAP activity was measured. Expression was tested in the following cell lines: Rodents- CHO (hamster) and C6 glioma, Primates-Vero, Human Hek 293 and Hela.

The results obtained indicate that in all the species tested, the addition of SP163 to the pBKC/S vector leads to a significant stimulation of expression of 5-10 fold as measured by the reporter SeAP. In the rat C6 glioma line SP163 addition leads to a 5-fold increase in SeAP activity (Figure 7), whereas in the Hela Cell line more than 10-fold stimulation was observed (Figure 7). The conclusion is that SP163 can be a useful addition to any vectors that are used to analyze expression of a target gene in transient transfection.

Effect of SP163 in stable expression in monocistronic vectors. Production of recombinant proteins for commercial purposes is done in systems where expression of the target gene is stable. The parental expression vector pBKC/S includes the neomycin gene that confers resistance to G418 (Figure 1). This feature was used to introduce the pBKC/S and pBKC/163S plasmids into CHO and C6 glioma cell lines. CHO is the preferred mammalian cell line for the production of biopharmaceuticals. Plasmid DNA of the vectors pBKC/S and pBKC163S was introduced by the liposome method and

colonies resistant to G418 were selected and analyzed. SeAP expression levels were tested in pools of the G418 resistant colonies. The results shown in Figure 8A-B and Fig. 11 demonstrate that the enhancing performance of SP163 is retained when the expression vector is integrated into the host cell genome. SP163 insertion in the 5' of the SeAP reporter gene led to a 5-20 fold stimulation of expression in CHO and C6 glioma cells.

The relative stimulation of SeAP activity (Figure 9), observed in plasmid pBKC/163S is not restricted to the comparison with plasmid pBKC/S. SeAP activity of pBKC/163S was compared in stable transfection in CHO cells with an additional two plasmids pCIBB/S to the basic expression vector and pCImc/S (pCIBB with the multiple cloning site that was engineered into pBKC to exclude the possibility that this segment is responsible for the enhancing effect). As shown in Figure 9, SeAP activity in pBKC/163S was four fold higher than pCIBB/S and over six-fold higher than pCImc/S.

EXAMPLE 5

SP163 ACTS AT THE LEVEL OF TRANSLATION

IN MONO-CISTRONIC AND BI-CISTRONIC VECTORS

SP163 enhances expression by increasing translation efficiency and not by influencing the abundance of mRNA due to increased stability or increased transcription. This was shown by Northern blot analysis of the amounts of mRNA of the reporter genes (SeAP and Luc) that are transcribed from the different vectors. The results are shown in Figure 10A-C.

The results for the mono-cistronic context are presented in Fig. 10A. The amounts of mRNA expressed by plasmids pBKC/S (control) [lane 1], pBKC/5'UTR/S [lane 2], pBKC/163/S [lane 3] and mock transfected cells [lane 4] were the same. The differences in the length of mRNA

generated by the different vectors reflect the addition in length due to the insert sizes (1 Kb in pBKC/5'UTR/S and 163 bp in pBKC/163/S).

5 The results for the bi-cistronic vector that are shown in Figure 10B and 10C indicate that the amounts of mRNA transcribed by the different cistrons are at the same level. Figure 10B shows results for the luciferase gene representing the first cistron and panel c for the second cistron SeAP (see Figure 1 for details). Each
10 cistron was analyzed with the corresponding probe by Northern blot analysis. The amounts of mRNA were the same for all plasmids: pBIC/LS (control) lane 1, pBIC/LUTRS lane 2, pBIC/L163S lane 3, pBIC/Lgrp78S (lane 5 is mock transfected cells).

15 As can be seen from the results in Figure 10B and C, only one mRNA is encoded by the bi-cistronic vectors without the IRES DNA (lane 1) or with the different DNA fragments tested (lane 2, 5'UTR, lane 3 SP163 and lane 4 grp78), clearly indicating that SP163 functions as an
20 IRES and that the stimulation of expression results from increased translation efficiency of the transcript and not from increase in the amount of the bi-cistronic mRNA or from the generation of a new SeAP mRNA that can be theoretically derived if SP163 has promoter activity.

25

EXAMPLE 6

TRANSLATIONAL ENHANCEMENT BY SP163

The possibility that the 5'UTR of VEGF (both the full length form and SP163) act as a translational
30 enhancer was examined by testing its ability to augment translation of a monocistronic mRNA. To this end, the full-size 5'UTR or SP163 was inserted between a CMV promoter and the coding region of SeAP in the context of a monocistronic vector (see Fig. 1). These constructs,
35 designated M/5'UTR and M/SP163 respectively, were

transfected into C6 cells and analyzed for SeAP production in pools of stably-transfected clones. Analysis of RNA extracted from transfected cells detected similar levels of SeAP-containing mRNA which was
5 represented in each case by a single band of the expected size (Fig. 11, inset). In contrast, the amount of SeAP activity released to the culture medium varied greatly according to the nature of the insert. Thus, the full-length 5'UTR augmented translation of SeAP by 5-fold and
10 SP163 enhanced SeAP translation much greater. In the experimental results shown in Fig. 11 enhancement by SP163 was 25-fold and in other experiments (not shown) enhancement was up to 40-fold.

Considerations discussed above led to the
15 examination of whether these elements also function as translational enhancers under conditions of hypoxic stress. As shown in Fig. 11, enhancement of SeAP translation was unaffected by growth under conditions of severe hypoxia.

20 To demonstrate that SP163 is a strong translational enhancer in additional cell types, the same constructs were also transfected into human 293 cells (Fig. 7) and hamster CHO cells (Fig. 8A). SP163 acted as a strong translation enhancer in these cells as well, augmenting
25 SeAP production to a level comparable to that shown for C6 cells (data not shown).

Throughout this application, various publications, are referenced by citation and patents by number. Full citations for the publications are listed below. The
30 disclosures of these publications and patents in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

The invention has been described in an illustrative
35 manner, and it is to be understood that the terminology

which has been used is intended to be in the nature of words of description rather than of limitation.

Obviously, many modifications and variations of the present invention are possible in light of the above
5 teachings. It is, therefore, to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described.

TABLE 1
SEQUENCE OF THE SP163 ELEMENT (SEQ ID No:7)

```

      10      20      30      ↓↓
AGCGCAGAGGCTTGGGGCAGCCGAGCGGCAGCCAGGCCC
40      50      60      70
CGGCCCCGGGCCTCGGTTCCAGAAGGGAGAGGAGCCCGCC
80      90      100     110
AAGGCGCGCAAGAGAGCGGGCTGCCTCGCAGTCCGAGCC
120     130     140     150
GGAGAGGGAGCGCGAGCCGCGCCCGGCCCCGGACGGCCTC-
160
CGAAACCATG

```

The oligonucleotide primers that were used for amplification of SP163 are underlined. These oligonucleotides correspond to the respective boundaries of the 1014 nucleotide long mouse VEGF 5'UTR. The initiation ATG codon is boxed. Arrows point at two possible locations of a presumptive splice junction. Nucleotide 1 of SP163 corresponds to nucleotides 1218 of the mouse VEGF gene, GeneBank accession U41383, while nucleotide 163 of the SP163 fragment corresponds to nucleotides 2231 of the mouse gene.

TABLE 2
BICISTRONIC AND MONOCISTRONIC CONSTRUCTS

<u>Bicistronic constructs</u>	<u>Insert</u>	<u>Monocistronic constructs</u>
B/0	NONE	M/0
B/UTR	1 VEGF 5' UTR 1014	M/UTR
B/SP163	1 SP163 163	M/SP163
B/BiP	1 BiP 5'UTR 221	(ND)

Designations of bicistronic and monocistronic constructs used in the Examples. The nucleotide numbers inside the rectangles represent the insert length. ND - not done.

REFERENCES

- Aiello, et al. (1994). Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other disorders [see comments]. N. Engl J Med 331:1480-7.
- Alon, et al. (1995). Vascular endothelial growth factor acts as a survival factor for newly formed retinal vessels and has implications for retinopathy of prematurity. Nat Med 1:1024-8.
- Banai, et al. (1994). Upregulation of vascular endothelial growth factor expression induced by myocardial ischemia: — implications for coronar angiogenesis. Cardiovasc Res 28:1176-9.
- Benda, et al. (1968). Differentiated rat glial cell strain in tissue culture. Science 161:370-74.
- Bernstein, et al. (1997). PDGF2/c-sis mRNA Leader Contains a Differentiation-linked Internal Ribosomal Entry Site (D-IRES). J. Biol. Chem. 272:9356-62.
- Carmeliet, et al. (1996). Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele. Nature 380:435-9.
- Chirgwin, et al. (1979). Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. Biochemistry 18:5294-99.
- Ehrenfeld, (1996). Initiation of Translation by Picornavirus RNAs. In Translational Control, J.W.B. Hershey, M.B. Mathews and N. Sonenberg, eds. (NY: Cold Spring Harbor Laboratory Press), pp. 549-73.
- Ferrara, et al. (1991). The vascular endothelial growth factor family of polypeptides. J. Cell Biochem 47:211-8.
- Ferrara, et al. (1996). Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. Nature 380:439-42.
- Finkenzellen, et al. (1995). Hypoxia-induced transcription of the vascular endothelial growth factor gene is independent of functional AP-1 transcription factor. Biochem. Biophys. Res. Comm. 208:432-439.

- Forsythe, et al. (1996). Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol* 16:4604-13.
- Frangakis and Kimelberg (1984). Dissociation of neonatal rat brain by dispase for the preparation of primary astrocyte culture. *Neurochem. Res.* 9:1689-98.
- Gan and Rhoads (1996). Internal Initiation of Translation Directed by the 5'-Untranslated Region of the mRNA for eIF4G, a factor Involved in the Picornavirus-induced Switch from Cap-dependent to Internal Initiation. *J. Biol. Chem.* 271:623-6.
- Heacock and Sutherland (1988). Induction and modulation of oxygen regulated protein synthesis. In *International Radiation Research Society Meeting* (Edinburgh, Scotland).
- Ikeda, et al (1995). Hypoxia-induced transcriptional activation and increased mRNA stability of vascular endothelial growth factor in C6 glioma cells. *J. Biol Chem* 270:19761-6.
- Jackson, (1996). A Comparative View of Initiation Site Selection mechanisms. In *Translational Control*, J.W.B. Hershey, M.B. Mathews and N. Sonenberg, eds. (NY: Cold Spring Harbor Laboratory Press), pp. 71-112.
- Jang, et al (1988). A segment of the 5' nontranslated region of encephalomyocarditisvirus RNA directs internal entry of ribosomes during in vitro translation. *J. Virol.* 62:2636-43.
- Jang, et al (1989). Initiation of Protein Synthesis by Internal Entry of Ribosomes into the 5' nontranslated Region of Encephalomyocarditis Virus RNA In Vivo. *J. Virol.* 63:1651-60.
- Jordan, et al. (1996). Transfecting mammalian cells: optimization of critical parameters affecting calcium-phosphate precipitate formation. *Nuc. Acid. Res.* 24:596-601.
- Kozak, (1991). An Analysis of Vertebrate mRNA Sequences: Intimations of Translational Control. *J. Cell. Bio.* 115:887-903.

Kraggerud, et al. (1995). Regulation of protein synthesis in human cells exposed to extreme hypoxia. *Anticancer Res.* 15:683-6.

Le and Maizel (1997). A common RNA structural motif involved in the internal initiation of translation of cellular mRNAs. *Nuc. Acid. Res.* 25:362-69.

Levy et al, 1995. Transcriptional regulation of the rat vascular endothelial growth factor gene by hypoxia. *J. Bio. Chem.* 270:13333-13340.

Lindquist (1986). The heat-shock response. *Ann. Rev. Biochem.* 55:1151-91.

Macejak and Sarnow (1991). Internal initiation of translation mediated by the 5' leader of a cellular mRNA. *Nature* 353:90-94.

Mathews et al (1996). Origins and targets of translational control. In *Translational Control*, J.W.B. Hershey, M.B. Mathews and N. Sonenberg, eds. (NY: Cold Spring Harbor Laboratory Press), pp. 1-29.

Matzura and Wennborg (1996). RNAdraw: an integrated program for RNA secondary structure calculation and analysis under 32-bit Microsoft Windows. *CABIOS* 12:247-9.

Mazure, et al. (1996). Oncogenic transformation and hypoxia synergistically act to modulate vascular endothelial growth factor expression. *Cancer Res* 56:3436-3340.

Meyuhas, et al. (1987). Glucocorticoids selectively inhibit translation of ribosomal protein and mRNAs in p1798 lymphosarcoma+++cells. *Mol. Cell. Biol.* 7:2691-99.

Meyuhas, et al. (1996). Translational control of ribosomal protein mRNAs in eukaryotes. In *Translational Control*, J.W.B. Hershey, M.B. Mathews and N. Sonenberg, eds. (NY: Cold Spring Harbor Laboratory Press), pp. 363-388.

Minchenko, et al. (1994). Hypoxic stimulation of vascular endothelial growth factor expression in vitro and in vivo. *Lab Invest.* 71:374-9.

Minty, et al. (1981). Mouse actin messenger RNAs. Construction and characterization of a recombinant plasmid molecule containing a complementary DNA transcript of mouse alpha-actin mRNA. J. Biol. 256:1008-14.

Mountford and Smith (1995). Internal ribosome entry sites and dicistronic RNAs in mammalian transgenesis. TIG 11(5):179-184.

Oh, et al. (1992). Homeotic gene Antennapedia mRNA contains 5'-noncoding sequences that confer translational initiation by internal ribosome binding. Genes. Dev. 6:1643-53.

Oh and Sarnow (1993). Gene Regulation: translational initiation by internal ribosome binding. Curr Opin Genetics Devel 3:295-300.

Pain (1996). Initiation of protein synthesis in eukaryotic cells. Eur. J. Biochem. 236:747-71.

Pelletier and Sonenberg (1988). Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA. Nature 334:320-325.

Pe'er, et al. (1995). Hypoxia-induced expression of VEGF by retinal cells is a common factor in neovascularizing ocular diseases. Lab. Invest. 72:638-645.

Shima et al., 1995. The mouse gene for vascular endothelial growth factor. J. Biol. Chem. 271:3877-3883.

Shweiki, et al. (1995). Induction of vascular endothelial growth factor expression by hypoxia and by glucose deficiency in multicell spheroids: implications for tumor angiogenesis. Proc. Natl. Acad. Sci. USA 92:768-72.

Sonenberg (1996). mRNA 5' Cap-binding Protein eIF4E and control of Cell Growth. In Translation Control, J.W. B. Hershey, M.B. Mathews and N. Sonenberg, eds. (NY: Cold Spring Harbor Laboratory Press), pp. 245-269.

Stein, et al. (1995). Stabilization of vascular endothelial growth factor mRNA by hypoxia and hypoglycemia and coregulation with other ischemia-induced genes. Mol. Cell. Biol. 15:5363-8.

Stone, et al. (1995). Development of retinal vasculature is mediated by hypoxia-induced vascular endothelial growth factor (VEGF) expression by neuralgia. *J Neurosci* 15:4738-47.

Teerink, et al. (1995). The human insulin-like growth factor II leader contains an internal ribosomal entry site. *Biochim. Biophys. Acta*. 1264:403-8.

Tischer, et al., 1991. The human gene for vascular endothelial growth factor. *J. Biol. Chem.* 266:11947-11954.

Vagner, et al. (1995). Alternative translation of human - fibroblast growth factor 2 mRNA occurs by internal entry of ribosomes. *Mol. Cell. Biol.* 15:35-44.

Ye, et al. (1997). Ultrabithorax and Antennapedia 5' Untranslated Regions Promote Developmentally Regulated Internal Translation Initiation. *Mol. Cell. Biol.* 17:1714-21.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Keshet, Eli
Itin, Ahuva
Stein, Ilan
- (ii) TITLE OF INVENTION: IRES SEQUENCES WITH HIGH TRANSLATIONAL
EFFICIENCY AND EXPRESSION VECTORS CONTAINING THE SEQUENCE
- (iii) NUMBER OF SEQUENCES: 7
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: KOHN & ASSOCIATES
 - (B) STREET: 30500 Northwestern Hwy. Suite 410
 - (C) CITY: Farmington Hills
 - (D) STATE: Michigan
 - (E) COUNTRY: US
 - (F) ZIP: 48334
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Montgomery, Ilene N.
 - (B) REGISTRATION NUMBER: 38,972
 - (C) REFERENCE/DOCKET NUMBER: 0168.00020
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (248) 539-5050
 - (B) TELEFAX: (248) 539-5055

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCGAGCGCA GAGGCTTGGG GC

22

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Primer"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CCATGGTTTC GGAGGCCGTC CG

22

- (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Primer"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTCGAGAGGT CGACGCCGGC CAAGACA

27

- (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Primer"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCATGGCTTG CCAGCCAGTT GGGCAGC

27

- (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Primer"

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GAGAGAATGA GCTTCCTACA G

21

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "Primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TCACCGCCTT GGCTTGTCAC A

21

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 166 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "SP163"
- (iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGCGCAGAGG CTTGGGGCAG CCGAGCGGCA GCCAGGCCCC GGCCCGGGCC TCGGTTCCAG	60
AAGGGAGAGG AGCCCGCCAA GCGCGCAAG AGAGCGGGCT GCCTCGCAGT CCGAGCCGGA	120
GAGGGAGCGC GAGCCGCGCC GGCCCGGAC GGCCTCCGAA ACCATG	166

CLAIMS

What is claimed is:

1. An isolated and cloned translation control element and analogues thereof having the nucleotide sequence as set forth in SEQ ID No:7 and designated SP163.

2. The translation control element as set forth in claim 1 wherein the sequence controls cap-independent mRNA translation via an internal ribosome entry site (IRES).

3. An expression vector comprising the translation control element as set forth in claim 1 operatively linked to a gene sequence to be expressed.

4. An expression vector comprising at least two nucleic acid sequences to be translated and said translation control sequence as set forth in claim 1 and wherein said translation control sequence is operatively linked to at least one of the sequences to be translated.

5. The expression vector as set forth in claim 4 wherein a single promoter is included.

6. A method for facilitating cap-independent translation of mRNA by including in an expression cassette a translation control element and analogues thereof having the nucleotide sequence as set forth in SEQ ID No:7 and designated SP163.

6. An internal ribosome entry site (IRES) having the nucleotide sequence as set forth in SEQ ID No:7.

7. A method for facilitating preferential translation of a gene of interest over the bulk of cellular mRNAs by including in an expression vector a translation control sequence and analogues thereof as set forth in SEQ ID No:7 operatively linked to the gene of interest and expressing the vector in host cells with reagents that inhibit cap-dependent translation.

8. A method for facilitating preferential translation of a gene of interest over the bulk of cellular mRNAs by including in an expression vector a translation control sequence and analogues thereof as set forth in SEQ ID No:7 operatively linked to the gene of interest and expressing the vector in host cells under conditions of cellular stress.




FIGURE 1A-B

2/11




FIGURE 2A-C

3/11





FIGURE 3A-B

4/11

**FIGURE 4**

5/11






FIGURE 5

6/11


**FIGURE 6**

7/11

**FIGURE 7**

8/11

A.



B.





FIGURE 8A-B

9/11

**FIGURE 9**

a.



1 2 3 4

b.



5 4 3 2 1


c.



5 4 3 2 1

FIGURE 10A-C

11/11

**FIGURE 11**

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/03699

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/11, 15/64, 15/67, 15/79

US CL : 435/320.1, 69.1, 172.3; 536/24.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 69.1, 172.3; 536/24.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG DATABASES: BIOSIS, MEDLINE, CA SEARCH, WORLD PATENT INDEX

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LE, S.-Y. et al. A common RNA structural motif involved in the internal initiation of translation of cellular mRNAs. Nucleic Acids Research. 15 January 1997, Vol. 25, No. 2, pages 362-369; see entire article, especially Figure 5.	1, 2
X	MORGAN, R. A. et al. Retroviral vectors containing putative internal ribosome entry sites: development of a polycistronic gene transfer system and applications to human gene therapy. Nucleic Acids Research. 25 March 1992, Vol. 20, No. 6, pages 1293-1299, see entire article.	1-6, 8



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
O documents referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

29 APRIL 1998

Date of mailing of the international search report

12 JUN 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHNNY F. RAILEY II, PH.D.

Telephone No. (703) 308-0196